

HEDIS Measure
<p>(FUH) Follow-Up After Hospitalization for Mental Illness</p> <p>Percentage of members with hospitalization for mental illness with a follow-up visit to behavioral health provider within 7 days of discharge</p> <p><i>*6-17 yr. old</i></p>
<p>(FUH) Follow-Up After Hospitalization for Mental Illness</p> <p>Percentage of members with hospitalizations for mental illness with a follow-up visit to behavioral health provider within 30 days of discharge</p> <p><i>*6-17 yr. old</i></p>
<p>(FUH) Follow-Up After Hospitalization for Mental Illness</p> <p>Percentage of members with hospitalizations for mental illness with a follow-up visit to behavioral health provider within 7 days of discharge</p> <p><i>*18 - 64 yr. old</i></p>
<p>(FUH) Follow-Up After Hospitalization for Mental Illness</p> <p>Percentage of members with hospitalizations for mental illness with a follow-up visit to behavioral health provider within 30 days of discharge</p> <p><i>*18-64 yr. old</i></p>
<p>(ADD) Follow-Up Care for Children Prescribed ADHD Medication</p> <p>Percentage of members with newly prescribed ADHD medication with 1 follow-up visit during the 30-day initiation phase</p> <p><i>*6-12 yr. old</i></p>
<p>(ADD) Follow-Up Care for Children Prescribed ADHD Medication</p> <p>Percentage of members with newly prescribed ADHD medication with at least 2 follow-up visits during the 10-month continuation and maintenance phase</p> <p><i>*6-12 yr. old</i></p>
<p>APC Children and Adolescents on 2 or more Antipsychotics</p> <p>Percentage of members on two or more concurrent antipsychotic medications (Lower rates are better)</p> <p><i>*12-17 yr. old</i></p>
<p>(AMM) Antidepressant Medication Management</p> <p>Effective Acute Phase Treatment: Percentage of members diagnosed with major depression who were treated with and remained on antidepressant medication for 12 weeks</p> <p><i>*18+ yr. old</i></p>
<p>(AMM) Antidepressant Medication Management</p>

<p>Effective Continuation Phase Treatment: Percentage of members diagnosed with major depression who were treated with and remained on antidepressant medication for 6 months</p> <p><i>*18+ yr. old</i></p>
<p>(SAA) Adherence to Antipsychotic Medications for Individuals With Schizophrenia</p> <p>Percentage of members with Schizophrenia who were dispensed and remained on antipsychotic medication for at least 80 percent of their treatment period.</p> <p><i>*18+ yr. old</i></p>
<p>(SSD) Diabetes Screening for People With Schizophrenia or Bipolar Disorder Who Are Using Antipsychotic Medications</p> <p>Percentage of members with Schizophrenia or Bipolar Disorder who were dispensed an antipsychotic medication and had a Diabetes screening test</p> <p><i>*18+ yr. old</i></p>
<p>(CDC) Comprehensive Diabetes Care</p> <p>Percentage of members with Diabetes who received an Hemoglobin A1c(HbA1c) test</p> <p><i>*18+ yr. old</i></p>
<p>(CDC) Comprehensive Diabetes Care</p> <p>Percentage of members with Diabetes whose HbA1c score indicated poor control (>9.0%)</p> <p><i>*18+ yr. old</i></p>
<p>(CDC) Comprehensive Diabetes Care</p> <p>Percentage of members with Diabetes whose HbA1c score indicated good control (< 8.0%)</p> <p><i>*18+ yr. old</i></p>
<p>(CDC) Comprehensive Diabetes Care</p> <p>Percentage of members with Diabetes who had an Eye exam (retinal) performed</p> <p><i>*18+ yr. old</i></p>
<p>(CDC) Comprehensive Diabetes Care</p> <p>Percentage of members with Diabetes who had medical attention for nephropathy</p> <p><i>*18+ yr. old</i></p>
<p>(CDC) Comprehensive Diabetes Care</p> <p>BP control (<140/90mm Hg)</p> <p><i>*18+ yr. old</i></p>
<p>(KED) Kidney Health Evaluation for Patients with Diabetes</p> <p>Percentage of members with diabetes who received a kidney health evaluation</p>

** 18-75 years of age*